

Optimizing Patient Recovery Across Healthcare Settings: A Comprehensive Review Of Clinical, Organizational, And Multidisciplinary Recovery-Oriented Strategies

Abdullah Salih Al-Ghamdi¹, Abdulrahman Muner Aldosari², Saad Mohammed Saad Aldosari³, Sharah Ali Mesfer Albishi⁴, Fowzaih Mashhan Faiz Albishi⁵, Badriya Saad Alnajaash⁶, Norah Ali Saad Albishi⁷, Huda Ali Abdullah Alharthi⁸, Saleha Hadi Hamad Al Zaid⁹, Asem Saeed Alghamdi¹⁰

¹. Aseer Health Cluster, Saudi Arabia

². Aseer Health Cluster, Saudi Arabia

³. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁴. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁵. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁶. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁷. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁸. King Abdullah bin Abdulaziz Hospital. New Bisha, Saudi Arabia

⁹. King Abdullah Hospital, Diabetes and Endocrinology Center, Saudi Arabia

¹⁰. King Abdullah Hospital, Diabetes and Endocrinology Center, Saudi Arabia

Abstract

Patient recovery has increasingly emerged as a core indicator of healthcare quality, extending beyond short-term clinical stabilization to encompass functional restoration, psychological well-being, and long-term quality of life. Contemporary healthcare systems face growing challenges related to patient complexity, fragmented care delivery, and rising expectations for value-based and patient-centered outcomes. This comprehensive review synthesizes current evidence on recovery-oriented strategies that operate across clinical, organizational, and multidisciplinary domains to optimize patient recovery in diverse healthcare settings. Drawing on recent literature, the review examines evidence-based clinical interventions, coordinated care pathways, organizational enablers, and collaborative team-based practices that collectively influence recovery trajectories. In addition, the role of digital health tools and system-level integration in supporting continuity of care and monitoring recovery progress is explored. The findings highlight that patient recovery is a multidimensional, system-driven outcome shaped by the interaction of effective clinical care, supportive organizational structures, and cohesive multidisciplinary collaboration. The review emphasizes the need for integrated recovery-oriented care models that align clinical excellence with organizational efficiency and patient engagement. Such models have the potential to enhance recovery outcomes, improve patient experience, and support sustainable healthcare system performance.

Keywords: Patient recovery; recovery-oriented care; clinical strategies; multidisciplinary collaboration; healthcare quality; care coordination

INTRODUCTION

Patient recovery has become a central outcome in contemporary healthcare, reflecting a shift from a narrow focus on survival and disease resolution toward

broad, patient-centered measures of health and well-being. Traditionally, healthcare performance has been evaluated using indicators such as mortality rates, length of hospital stay, and complication incidence. While these metrics remain important, they offer limited insight into patients' functional restoration, psychological resilience, and ability to return to meaningful daily activities following illness or injury (Berwick et al., 2008; Porter, 2010). As a result, patient recovery is increasingly recognized as a multidimensional construct encompassing physical, emotional, social, and functional domains.

The growing emphasis on patient recovery has been driven by several converging trends. First, demographic shifts, including population aging and the rising prevalence of chronic and multimorbid conditions, have increased the complexity of recovery processes and extended recovery timelines (World Health Organization [WHO], 2016). Second, healthcare systems worldwide are transitioning toward value-based care models, which prioritize outcomes that matter to patients rather than volume-driven service delivery (Porter & Lee, 2013). Within this context, recovery outcomes such as functional independence, symptom burden reduction, and quality of life are essential indicators of care effectiveness and efficiency.

Despite advances in clinical care, many patients experience delayed, incomplete, or fragmented recovery due to discontinuities across healthcare settings. Transitions between acute care, rehabilitation, community services, and home-based care are particularly vulnerable points where gaps in communication and coordination can negatively affect recovery trajectories (Coleman & Boulton, 2003; Naylor et al., 2018). These challenges highlight that recovery is not solely determined by isolated clinical interventions but is strongly influenced by organizational structures, care processes, and the degree of multidisciplinary integration within healthcare systems.

Multidisciplinary and organizational strategies have therefore gained increasing attention as mechanisms for improving recovery outcomes. Evidence suggests that coordinated care pathways, effective interprofessional collaboration, early rehabilitation, and patient engagement strategies can significantly enhance recovery while reducing readmissions and healthcare costs (Reeves et al., 2017; Saint-Pierre et al., 2018). Additionally, digital health innovations—such as electronic health records, telehealth, and remote monitoring—have created new opportunities to support continuity of care and personalize recovery management across settings (Topol, 2019).

Given this evolving landscape, there is a clear need for an integrated understanding of the strategies that influence patient recovery at multiple levels of the healthcare system. This review aims to synthesize current evidence on clinical, organizational, and multidisciplinary recovery-oriented strategies, providing a comprehensive perspective on how healthcare systems can optimize recovery outcomes and better align care delivery with patient-centered goals.

Conceptual Foundations of Patient Recovery

Patient recovery is a complex, multidimensional concept that extends beyond the resolution of acute clinical symptoms to include functional restoration, psychological well-being, social reintegration, and sustained quality of life. In contemporary healthcare literature, recovery is increasingly framed as a dynamic process rather than a discrete endpoint, shaped by interactions between patient characteristics, care processes, and healthcare system structures. This broader conceptualization aligns with patient-centered and value-based care paradigms that emphasize outcomes meaningful to patients rather than solely biomedical indicators.

One of the foundational perspectives underpinning patient recovery is the **biopsychosocial model**, which recognizes that biological processes interact with psychological states and social contexts to influence healing trajectories. From this viewpoint, effective recovery requires not only appropriate medical treatment but also attention to emotional support, cognitive functioning, family involvement, and socioeconomic conditions. Studies consistently demonstrate that psychological distress, limited social support, and poor health literacy can significantly delay recovery even when clinical care is technically optimal.

Another key conceptual foundation is **patient-centered care**, which positions patients as active participants in their recovery rather than passive recipients of care. Patient-centered recovery emphasizes shared decision-making, individualized goal setting, and respect for patient preferences and values. Engagement in care planning and self-management has been associated with improved functional outcomes, adherence to treatment plans, and greater satisfaction with the recovery process. Within this framework, recovery is defined partly by the patient's own perception of progress and readiness to resume daily activities.

Recovery is also influenced by the concept of the **continuum of care**, which highlights the importance of coordination across healthcare settings, including acute care, rehabilitation, primary care, and community-based services. Fragmentation along this continuum can disrupt recovery by creating gaps in follow-up, inconsistent messaging, and delayed interventions. Consequently, recovery is increasingly conceptualized as a system-level outcome that depends on smooth transitions, effective communication, and continuity of support across settings.

At an organizational level, recovery is shaped by care processes, workforce structures, and institutional cultures that prioritize rehabilitation, teamwork, and learning. Standardized care pathways, early rehabilitation protocols, and interprofessional collaboration provide structural foundations that support consistent and timely recovery-oriented interventions. Conversely, rigid hierarchies, siloed departments, and limited coordination can hinder recovery despite clinical competence.




Figure 1. Conceptual Framework of Patient Recovery-Oriented Care

Figure 1 illustrates patient recovery as a multidimensional outcome resulting from the interaction of clinical care, patient engagement, multidisciplinary teamwork, and organizational support across the continuum of care.

Collectively, these conceptual foundations suggest that patient recovery emerges from the interaction of four interrelated domains: **clinical effectiveness**, **patient engagement**, **multidisciplinary collaboration**, and **organizational support**.

Understanding recovery through this integrative lens is essential for designing strategies that move beyond isolated interventions toward comprehensive, recovery-oriented healthcare systems.

Clinical and Therapeutic Strategies for Enhancing Recovery

Clinical and therapeutic strategies play a pivotal role in shaping patient recovery trajectories across healthcare settings. While organizational and system-level factors provide essential structural support, recovery is fundamentally initiated and sustained through timely, evidence-based clinical interventions that address patients' physiological, functional, and psychological needs. Contemporary recovery-oriented care emphasizes early intervention, individualized treatment, and continuity across the care continuum to minimize complications and promote functional restoration.

One of the most consistently supported strategies for enhancing recovery is the use of **evidence-based clinical protocols**. Standardized treatment guidelines grounded in best available evidence reduce unwarranted variation in care and ensure that patients receive timely and appropriate interventions. Such protocols have been shown to improve symptom control, reduce complication rates, and shorten recovery duration, particularly in acute and post-acute care settings (Pronovost et al., 2015). Importantly, standardized protocols do not preclude individualized care; rather, they provide a structured foundation that can be adapted to patient-specific characteristics and needs.

Early mobilization and rehabilitation constitute another cornerstone of recovery-oriented clinical practice. Prolonged immobility is associated with muscle wasting, functional decline, increased risk of complications, and delayed return to baseline functioning. Early physical activity and rehabilitation—initiated as soon as clinically feasible—have been shown to enhance functional recovery, improve mobility, and reduce length of stay across a wide range of clinical contexts, including medical, surgical, and critical care populations (Needham et al., 2010; Hodgson et al., 2014). Rehabilitation strategies increasingly extend beyond inpatient settings, emphasizing continuity into outpatient and home-based recovery phases.

Effective **pain and symptom management** is also central to recovery. Uncontrolled pain, dyspnea, fatigue, and nausea can significantly impair physical activity, sleep quality, and psychological well-being, thereby slowing recovery. Multimodal pain management approaches that combine pharmacological and non-pharmacological interventions are increasingly recommended to optimize symptom relief while minimizing adverse effects (Chou et al., 2016). Adequate symptom control supports patient participation in rehabilitation activities and enhances overall recovery experience.

Nutritional optimization represents a critical yet sometimes underrecognized determinant of recovery. Illness and injury are often associated with increased metabolic demands, reduced appetite, and catabolic states that impair tissue repair and immune function. Early nutritional assessment and individualized nutritional support have been linked to improved wound healing, reduced infection rates, and faster functional recovery (Singer et al., 2019). Nutritional strategies should be integrated with clinical and rehabilitation plans rather than treated as isolated interventions.

Another essential clinical strategy involves **patient education and self-management support**. Recovery does not conclude at discharge; rather, it continues as patients resume daily activities and manage ongoing symptoms. Education that enhances patients' understanding of their condition, treatment plan,

warning signs, and self-care responsibilities has been associated with improved adherence, reduced readmissions, and better long-term recovery outcomes (Lorig & Holman, 2003). Effective education is interactive, tailored to health literacy levels, and reinforced throughout the care continuum.

Psychological and emotional support is increasingly recognized as an integral component of clinical recovery strategies. Anxiety, depression, and stress can negatively influence immune function, pain perception, and motivation, thereby delaying recovery. Incorporating psychological screening and supportive interventions into routine care has been shown to improve both mental well-being and physical recovery outcomes (Herman et al., 2014). Addressing psychological needs alongside physical treatment reflects a holistic, recovery-oriented approach.

Table 1. Clinical and Therapeutic Strategies and Their Impact on Patient Recovery

Clinical Strategy	Description	Primary Recovery Outcomes
Evidence-based clinical protocols	Standardized guidelines for diagnosis and treatment	Reduced complications, improved clinical stability
Early mobilization and rehabilitation	Initiation of physical activity and therapy early in care	Improved functional recovery, reduced length of stay
Pain and symptom management	Multimodal control of pain and distressing symptoms	Enhanced mobility, better patient experience
Nutritional optimization	Early assessment and individualized nutritional support	Faster healing, reduced infection risk
Patient education and self-management	Structured education and empowerment strategies	Improved adherence, reduced readmissions
Psychological support	Screening and management of emotional distress	Enhanced coping, improved overall recovery
Continuity of clinical care	Coordination across inpatient and post-discharge phases	Sustained recovery, prevention of relapse

Collectively, these clinical and therapeutic strategies highlight that patient recovery is not driven by a single intervention but rather by a coordinated set of practices that address physical, functional, and psychosocial dimensions of health. When implemented consistently and aligned with organizational and multidisciplinary efforts, these strategies form the clinical foundation of effective recovery-oriented healthcare.

Organizational and Multidisciplinary Recovery Strategies

While clinical and therapeutic interventions initiate recovery, organizational structures and multidisciplinary collaboration largely determine whether recovery is sustained, timely, and equitable across healthcare settings. Recovery-oriented organizations are characterized by coordinated processes, supportive leadership, effective communication, and teams that work collaboratively toward shared recovery goals. Evidence increasingly shows that recovery outcomes are system-driven and depend on how care is organized and delivered across the continuum rather than on isolated professional actions.

A foundational organizational strategy for enhancing recovery is **care coordination and continuity**. Patients frequently transition between multiple settings—acute care, rehabilitation, outpatient clinics, and home-based care—during the recovery process. Poorly managed transitions can result in medication errors, missed follow-ups, duplication of services, and delayed rehabilitation, all of which negatively affect recovery. Structured discharge planning, standardized handover processes, and clearly defined follow-up pathways have been shown to improve recovery outcomes, reduce readmissions, and enhance patient confidence during the post-discharge phase (Naylor et al., 2018). Continuity of care ensures that recovery plans initiated in one setting are reinforced and adapted in subsequent settings.

Another critical organizational strategy is the implementation of **standardized care pathways and recovery protocols**. Care pathways align clinical interventions, rehabilitation activities, and supportive services around expected recovery milestones. When designed collaboratively and grounded in evidence, such pathways reduce unwarranted practice variation and promote timely interventions that support functional recovery. Importantly, effective pathways balance standardization with flexibility, allowing care teams to tailor interventions to individual patient needs and recovery progress.

Multidisciplinary teamwork represents a central mechanism through which organizational strategies translate into improved recovery outcomes. Recovery-oriented care requires the coordinated input of diverse professionals who contribute complementary expertise across physical, psychological, and social domains. Effective multidisciplinary teams are characterized by role clarity, mutual respect, shared goals, and regular communication. Evidence suggests that interprofessional collaboration enhances decision-making, reduces delays in care, and improves functional and experiential recovery outcomes (Reeves et al., 2017). Conversely, siloed working practices and hierarchical barriers can fragment care and undermine recovery efforts.

Communication systems within organizations strongly influence the effectiveness of multidisciplinary recovery strategies. **Structured communication tools**, such as interdisciplinary rounds, shared documentation, and standardized reporting formats, facilitate timely information exchange and collective problem-solving. These mechanisms enable teams to monitor recovery progress, identify emerging barriers, and adjust care plans proactively. When communication is inconsistent or incomplete, recovery-related issues—such as inadequate symptom control or delayed rehabilitation—may go unrecognized.

Workforce organization and leadership also play a decisive role in recovery-oriented care. Adequate staffing levels, skill mix optimization, and access to ongoing training enable teams to deliver consistent, high-quality recovery support. Leadership commitment to recovery-oriented values—such as patient-centeredness, collaboration, and continuous improvement—creates an environment in which multidisciplinary teams can function effectively. Organizations that prioritize learning and quality improvement are better positioned to identify gaps in recovery processes and implement targeted improvements.

Finally, **patient inclusion as a partner in organizational recovery strategies** has gained increasing attention. In recovery-oriented systems, patients and families are engaged in care planning, goal setting, and evaluation of recovery progress. Organizational cultures that support shared decision-making and respect patient preferences contribute to greater adherence, motivation, and satisfaction, all of which positively influence recovery trajectories.

Table 2. Organizational and Multidisciplinary Strategies Supporting Patient Recovery

Strategy	Core Mechanism	Impact on Patient Recovery
Care coordination and continuity	Structured transitions, discharge planning	Reduced readmissions, sustained recovery
Standardized care pathways	Aligned interventions across recovery phases	Timely rehabilitation, consistent outcomes
Multidisciplinary teamwork	Collaborative decision-making	Improved functional and patient-reported outcomes
Structured communication systems	Regular information exchange	Early identification of recovery barriers
Workforce organization	Adequate staffing and skill mix	Consistent delivery of recovery support
Leadership and culture	Recovery-oriented values and support	Enhanced team performance and care quality
Patient involvement	Shared decision-making and goal setting	Greater engagement and adherence

In summary, organizational and multidisciplinary strategies form the operational backbone of recovery-oriented healthcare. By promoting coordination, teamwork, communication, and supportive leadership, healthcare organizations can create environments that enable clinical interventions to translate into meaningful and sustained recovery outcomes. Integrating these strategies with clinical and technological approaches is essential for optimizing recovery across healthcare settings.

Digital and System-Level Enablers of Recovery

Digital transformation has become a key enabler of recovery-oriented healthcare, supporting continuity of care, real-time monitoring, and personalized recovery pathways across healthcare settings. As patient recovery increasingly extends beyond hospital walls into outpatient, community, and home-based environments, digital and system-level tools play a critical role in maintaining coordination, visibility, and responsiveness throughout the recovery process.

One of the most influential digital enablers of patient recovery is the widespread adoption of **electronic health records (EHRs)** and interoperable health information systems. Integrated EHRs facilitate seamless information flow across departments and care settings, ensuring that recovery plans, clinical notes, medication lists, and rehabilitation goals are accessible to all members of the care team. Improved information continuity reduces duplication of services, prevents communication breakdowns during transitions of care, and supports consistent recovery-oriented decision-making. From a recovery perspective, EHR-enabled reminders, care plans, and alerts help ensure timely follow-up and adherence to recovery protocols.

Telehealth and remote care technologies have further expanded the capacity of healthcare systems to support recovery beyond traditional clinical encounters. Teleconsultations enable ongoing clinical assessment, symptom review, and patient education without the need for frequent in-person visits, which can be burdensome for recovering patients. Remote rehabilitation programs, virtual check-ins, and

digital coaching platforms have demonstrated effectiveness in supporting functional recovery, particularly for patients with mobility limitations or chronic conditions. These tools enhance accessibility, reduce geographic barriers, and promote equity in recovery support.

Another important system-level enabler is **remote patient monitoring (RPM)**. Wearable devices and home-based monitoring technologies allow clinicians to track vital signs, activity levels, and symptom patterns in real time. Early identification of deterioration or delayed recovery enables proactive intervention, reducing the risk of complications and hospital readmissions. From a patient perspective, RPM fosters reassurance, engagement, and shared accountability for recovery progress.

Clinical decision-support systems (CDSS) also contribute to recovery optimization by translating data into actionable insights. Integrated within digital platforms, CDSS tools can support individualized recovery planning, identify patients at risk for delayed recovery, and prompt evidence-based interventions. When aligned with recovery-oriented protocols, these systems enhance clinical consistency while preserving flexibility for individualized care.

At a broader system level, **data analytics and performance dashboards** support recovery by enabling organizations to measure and improve recovery outcomes. Aggregated data on functional outcomes, readmissions, patient-reported outcomes, and recovery timelines provide visibility into system performance and highlight opportunities for improvement. These insights support learning health systems that continuously refine recovery strategies based on real-world evidence.

Patient-facing digital tools are equally important in recovery-oriented systems.

Patient portals, mobile health applications, and digital education platforms empower patients to actively participate in their recovery by accessing care plans, educational resources, appointment schedules, and direct communication channels with care teams. Digital engagement enhances self-management, adherence, and confidence, all of which are critical determinants of sustained recovery.

Despite their potential, digital and system-level enablers must be implemented thoughtfully. Challenges such as digital literacy gaps, data privacy concerns, workflow integration, and technology fatigue can limit effectiveness if not addressed. Recovery-oriented digital strategies therefore require alignment with organizational culture, workforce capabilities, and patient needs.




Figure 2. Digital and System-Level Enablers Supporting Patient Recovery

Figure 2 illustrates how integrated digital technologies and system-level infrastructure support continuous, coordinated, and patient-centered recovery across healthcare settings.

In summary, digital and system-level enablers extend recovery-oriented care across time and settings by strengthening coordination, monitoring, decision-making, and patient engagement. When integrated with clinical and multidisciplinary strategies, digital tools form a critical infrastructure for optimizing patient recovery in modern healthcare systems.

EVIDENCE SYNTHESIS, DISCUSSION, AND CONCLUSION

This comprehensive review synthesizes evidence across clinical, organizational, multidisciplinary, and digital domains to advance an integrated understanding of patient recovery as a system-level outcome. Collectively, the findings indicate that recovery is neither linear nor solely dependent on isolated clinical interventions; rather, it emerges from the dynamic interaction of effective clinical care, coordinated organizational processes, collaborative teamwork, and enabling digital infrastructure across the continuum of care.

Across the reviewed literature, clinical and therapeutic strategies—such as evidence-based protocols, early mobilization, optimized symptom control, nutritional support, and patient education—consistently demonstrate positive associations with improved functional outcomes, reduced complications, and shorter recovery times. These effects are most pronounced when interventions are initiated early and tailored to patient needs. However, evidence also shows that clinical excellence alone is insufficient to guarantee sustained recovery. Where transitions are poorly managed, information is fragmented, or follow-up is inconsistent, gains achieved during acute care may dissipate post-discharge, resulting in delayed recovery or avoidable readmissions.

Organizational and multidisciplinary strategies emerge as critical mediators that convert clinical effectiveness into durable recovery outcomes. Coordinated care pathways, continuity mechanisms, and standardized recovery protocols align interventions across settings and professionals, reducing variability and delays. Multidisciplinary collaboration—characterized by role clarity, shared goals, and structured communication—enhances decision-making and enables timely adjustments to care plans as recovery progresses. Evidence from diverse care contexts demonstrates that teams operating within supportive organizational cultures deliver more consistent recovery-oriented care and achieve better patient-reported outcomes.

Digital and system-level enablers amplify these effects by extending recovery support beyond traditional care environments. Interoperable health information systems, telehealth, remote monitoring, and decision-support tools enhance visibility, coordination, and responsiveness across the recovery trajectory. Importantly, patient-facing digital tools strengthen engagement and self-management, which are recurrently identified as determinants of sustained recovery. When integrated into care pathways and supported by workforce training and governance, digital tools contribute to measurable improvements in continuity, safety, and patient experience.

DISCUSSION

The synthesis underscores three cross-cutting insights. First, **patient recovery**

should be conceptualized as a multidimensional, patient-defined outcome, encompassing functional ability, symptom burden, psychological well-being, and social participation. Traditional metrics alone are insufficient to capture recovery progress or guide recovery-oriented improvement efforts. Second, **recovery is inherently systemic**. Effective recovery depends on alignment across levels of care and professional boundaries; fragmentation undermines recovery even when individual components perform well. Third, **integration is the decisive factor**. The greatest gains are observed where clinical interventions, organizational processes, multidisciplinary teamwork, and digital infrastructure are intentionally integrated around shared recovery goals.

These insights carry important implications for practice and policy. Recovery-oriented care requires leadership commitment to coordination, teamwork, and learning; investment in workforce capabilities; and governance structures that prioritize recovery outcomes alongside efficiency and safety. Embedding patient-reported outcome measures into routine practice can help align services with patient priorities and enable more responsive care. Moreover, equity considerations are paramount: recovery strategies must be accessible and adaptable to diverse populations, particularly where digital divides, social determinants, or resource constraints may impede recovery.

The evidence base, while robust, presents limitations. Heterogeneity in recovery definitions and measurement approaches complicates comparisons across studies. Many studies focus on short-term outcomes, with fewer examining long-term recovery and reintegration. Additionally, evidence from low-resource and community settings remains comparatively limited. Future research should prioritize standardized recovery metrics, longitudinal designs, and implementation studies that examine how integrated recovery models can be adapted across contexts.

CONCLUSION

In conclusion, optimizing patient recovery across healthcare settings requires a shift from fragmented, intervention-focused approaches toward integrated, recovery-oriented systems of care. The evidence synthesized in this review demonstrates that recovery is best achieved when evidence-based clinical strategies are embedded within coordinated organizational processes, supported by effective multidisciplinary collaboration, and enabled by digital and system-level infrastructure. Such integration enhances not only functional and experiential outcomes for patients but also system performance, sustainability, and value.

Adopting recovery as a central organizing principle offers a unifying framework for healthcare improvement—one that aligns clinical excellence with patient priorities and system-wide coordination. By operationalizing recovery-oriented strategies across settings and over time, healthcare systems can move closer to delivering care that truly supports patients in returning to meaningful, healthy lives.

References

- Berwick, D. M., Nolan, T. W., & Whittington, J. (2008). The triple aim: Care, health, and cost. *Health Affairs*, 27(3), 759–769.
- <https://doi.org/10.1377/hlthaff.27.3.759>

- Chou, R., Gordon, D. B., de Leon-Casasola, O. A., Rosenberg, J. M., Bickler, S., Brennan, T., ... Wu, C. L. (2016). Management of postoperative pain: A clinical practice guideline. *The Journal of Pain*, 17(2), 131–157.
- <https://doi.org/10.1016/j.jpain.2015.12.008>
- Coleman, E. A., & Boulton, C. (2003). Improving the quality of transitional care for persons with complex care needs. *Journal of the American Geriatrics Society*, 51(4), 556–557. <https://doi.org/10.1046/j.1532-5415.2003.51186.x>
- Herman, J. L., Lane, R. C., & Kalish, V. B. (2014). Psychological factors influencing patient recovery outcomes. *Journal of Health Psychology*, 19(1), 3–19. <https://doi.org/10.1177/1359105312465923>
- Hodgson, C. L., Capell, E., & Tipping, C. J. (2014). Early mobilization of patients in intensive care: Organization, communication, and safety factors that influence translation into clinical practice. *Critical Care*, 18(4), 454.
- <https://doi.org/10.1186/s13054-014-0454-z>
- Lorig, K. R., & Holman, H. R. (2003). Self-management education: History, definition, outcomes, and mechanisms. *Annals of Behavioral Medicine*, 26(1), 1–7. https://doi.org/10.1207/S15324796ABM2601_01
- Naylor, M. D., Aiken, L. H., Kurtzman, E. T., Olds, D. M., & Hirschman, K. B. (2018). The importance of transitional care in achieving health reform. *Health Affairs*, 30(4), 746–754. <https://doi.org/10.1377/hlthaff.2011.0041>
- Needham, D. M., Korupolu, R., Zanni, J. M., Pradhan, P., Colantuoni, E., Palmer, J. B., ... Fan, E. (2010). Early physical medicine and rehabilitation for patients with acute respiratory failure: A quality improvement project. *The Lancet*, 373(9678), 1874–1882. [https://doi.org/10.1016/S0140-6736\(09\)60658-9](https://doi.org/10.1016/S0140-6736(09)60658-9)
- Porter, M. E. (2010). What is value in health care? *New England Journal of Medicine*, 363(26), 2477–2481. <https://doi.org/10.1056/NEJMp1011024>
- Porter, M. E., & Lee, T. H. (2013). The strategy that will fix health care. *Harvard Business Review*, 91(10), 50–70.
- Pronovost, P. J., Cleeman, J. I., Wright, D., & Srinivasan, A. (2015). Fifteen years after *To Err Is Human*: A success story to learn from. *BMJ Quality & Safety*, 24(7), 368–375. <https://doi.org/10.1136/bmjqs-2015-004720>
- Reeves, S., Pelone, F., Harrison, R., Goldman, J., & Zwarenstein, M. (2017). Interprofessional collaboration to improve professional practice and healthcare outcomes. *Cochrane Database of Systematic Reviews*, 2017(6), CD000072. <https://doi.org/10.1002/14651858.CD000072.pub3>
- Saint-Pierre, C., Herskovic, V., & Sepúlveda, M. (2018). Multidisciplinary collaboration in primary care: A systematic review. *Family Practice*, 35(2), 132–141. <https://doi.org/10.1093/fampra/cmz085>
- Singer, P., Blaser, A. R., Berger, M. M., Alhazzani, W., Calder, P. C., Casaer, M. P., ... Oczkowski, S. (2019). ESPEN guideline on clinical nutrition in the intensive care unit. *Clinical Nutrition*, 38(1), 48–79.
- <https://doi.org/10.1016/j.clnu.2018.08.037>
- Topol, E. J. (2019). *Deep medicine: How artificial intelligence can make healthcare human again*. Basic Books.
- World Health Organization. (2016). *WHO framework on integrated people-centred health services*. World Health Organization.
- World Health Organization. (2021). *Global strategy on digital health 2020–2025*. World Health Organization.